

## AUTHOR KORESPONDENSI

Judul Artikel : Whether Beet Juice and Red Dragon Fruit Juice Increase the Haemoglobin Level in Young People?


**Identitas Proceeding** : Proceedings of the 1st Borobudur International Symposium on Humanities, Economics and Social Sciences (BIS-HESS 2019)

- a. Nama Proceeding : atlantis - press
- b. Nomor ISSN : 2352-5398
- c. Volume, Nomor, Bulan Tahun : 4 Mei 2020
- d. Penerbit : Atlantis Press.
- e. DOI artikel (jika ada) : [10.2991/assehr.k.200529.134](https://doi.org/10.2991/assehr.k.200529.134)
- f. Alamat Web Jurnal : <https://www.atlantis-press.com/proceedings/bis-hess-19/125939583>
- g. Terindeks di : Web of Science, Scopus, MEDLINE, PubMed Central (PMC), DOAJ, Ei Compendex, PubMed, Portico, Ulrichsweb, Google scholar

### Riwayat Artikel

1. Manuscript submitted : 6 Agustus 2019
2. Manuscript revised : 27 September 2019
3. Accepted for publication : 28 Oktober 2019
4. Published : 4 Mei 2020

### Lampiran Pendukung :



[Print this page](#)

# BIS 2019

1st Borobudur International Symposium 2019  
Grand Artos Hotel and Convention - Magelang, 16 October 2019  
Website: <http://bis.ummgl.ac.id>  
Email: [dppj@ummgl.ac.id](mailto:dppj@ummgl.ac.id)

---

Date: 15 August 2019

## Letter of Acceptance

Dear Authors: Endang Wahyuningsih, Endang Sawitri, Choiril Hana

We are pleased to inform you that your full paper (ABS-448, Oral Presentation), entitled:

**"Whether Beet Juice and Red Dragon Fruit Juice Increase the  
Haemoglobin Level in Young People?"**

has been reviewed and accepted to be published at Advances in Social Science, Education and Humanities Research Altantis Press.

Please visit our website for more information.

Thank You.

Best regards,

Dr. Muji Setiyo, ST.,MT  
BIS 2019 Chairperson

# THE INNOVATION APPLICATION OF BEET JUICE AND RED DRAGON FRUIT JUICE TO INCREASING HAEMOGOBBLIN LEVEL IN YOUNG PEOPLE

EndangWahyuningsih<sup>1</sup>, EndangSawitri<sup>1</sup>, Choiril Hana<sup>1</sup>

<sup>1</sup>Midwifery Department, Sekolah Tinggi Ilmu Kesehatan Muhammadiyah

<sup>2</sup>Nursing Department, Sekolah Tinggi Ilmu Kesehatan Muhammadiyah

<sup>3</sup>Pharmacy Department, Sekolah Tinggi Ilmu Kesehatan Muhammadiyah

\*endangwahyuningsih@stikesmukla.ac.id

**Abstract.** Anaemia has been one of the social health problems which arises and is spread around the world especially in developing countries and poor countries. Anaemia affects a large number of people mainly young males and females. Women and young females need twice as much iron as men and young males do. WHO states that 25 – 40% of young people suffer from anaemia and it strikes more than 57% of young females in Indonesia. The red beet is rich in various vitamin B content: vitamin B1, B2, B3 and B6. The dragon fruit also contains iron, vitamin B1, vitamin B2, vitamin B and vitamin C. Our research purpose is to find out the comparison of beet juice and dragon fruit juice consumption to increasing haemoglobin level in young people. The research design is *Quasi-Experiment* using *pre test-post test with control group design*. The technique sampling we use is *purposive sampling*. We use a *juice extractor* and *haemometer* to study the samples for seven days. There were 32 samples in our research. Our research used paired t-test and independent t-test. The result showed that beet *p value* was  $0,515 > 0,05$  and dragon fruit was  $0,516 > 0,05$ , and that the consumption of beet juice and dragon fruit juice did not affect the haemoglobin level in young females.

## 1. Introduction

Young people still show growth especially their reproductive system. During menstrual period, females need enough iron. Protein, iron and vitamin C intakes are related to menstrual cycle [1]. Dysmenorrhoea is an annoying problem for some women, especially for those who do intense activities like young people. One of the factors which affect dysmenorrhoea is the Hb level is less than 12gr/% (anaemia) which often strikes young people. The main factor which contributes to less haemoglobin level is bleeding during menstrual period and insufficient nutrients intake. Iron and proteins young people take will reduce dysmenorrhoea and anaemia [2]. Young females face a high risk of suffering from anaemia because they have a menstrual period every month. They can increase their haemoglobin level by take vitamin C and tablet Fe [3]. [4] Those who suffer from anaemia lack Fe, Zn, Cu, folic acid and B6 intake. Lack vitamin A and C will also affect their haemoglobin level [5]. [6] Vegetable protein also plays an important role in anaemia development in WUS. [7] The young females' haemoglobin level who live in mountainous regions is different from that of who live in coastal regions. [8] Children who suffer from GAKI (a condition in which there is a deficiency of iodine) in Indonesia are related to anaemia. Eating habit will affect the size of anthropometry including weight, height and abdominal circumference of young people's. Eating habit in which there is a deficiency of nutrient will affect the abnormality of anthropometry size which results in anaemia [9]. Young people's unhealthy life style such as consuming fast food, drinking soda, drinking unhealthy energy drinks will result in health problems and chronic diseases which are not contagious so that it increases mortality and morbidity [10]. In assosa zone, anaemia occurs in socio-demographic conditions and food intake [11]. According to the data from Muhammad Al Kaustar Junior High School, there are 235 teenagers aged 13-16. Based on the information from student department, a clinic in Kartosuro has a counseling program to measure the students' weight and height every three month. Apart from that, there has never been anaemia skimming as an early detection of anaemia in

young people conducted by the school and clinic in Kartosuro. This will affect their concentration and achievements.

## 2. Methods

Our research design used *quasi experimental* with research design being used was *non equivalent pre and post test control group design*. We separated the group into two groups. The first group was given beet juice and the second was given red dragon fruit juice. We checked the young females' haemoglobin level. After we collected the data, we took the respondents based on their criteria. The result from the skimming of their haemoglobin level showed that there were 32 people with abnormal haemoglobin level. After we collected the data, the respondents' data was grouped into 2 groups. In the first group, 16 respondents were given beet juice, and in the second group 16 respondents were given red dragon fruit juice. They were beet juice without water as much as 100 grams for 7 days by using *a juice extractor*. The second group were given red dragon fruit juice without water as much as 100 grams for 7 days by using *a juice extractor*. Then we conducted the *post test* to measure the first and second's group haemoglobin level using *paired t-test* and *independent t-test*. The *paired t-test* was conducted to test the different mean of the two results of measurement in the same group. While the *independent t-test* was conducted to test the different mean of both groups of interventions/cases and the control groups.

## 3. Result and Discussion

Based on the research results, it was known that the average haemoglobin level before the pre test (before they were given beet juice) was lower which was 10.94 gr/dl compared to the average haemoglobin level after the post test (after they were given beet juice) which was 11.58 gr/dl. With the highest score of pre test haemoglobin level (11.3 gr/dl), it was lower compared to the post test (13.6 gr/dl), and the lowest score of pre test was 6.7 gr/dl was lower compared to the haemoglobin level after they were given beet juice which was 9.9 gr/dl. This proved that the average haemoglobin level of pre test after beet juice was given was lower than that of post test (10.49 gr/dl < 11.58 gr/dl).

Based on the research results, the average haemoglobin level of pre test (before they were given red dragon fruit juice) showed lower number which was 11.63 ge/dl compared to the average haemoglobin level of post test (after they were given red dragon fruit juice) which was 12.36 gr/dl. The haemoglobin level of pre test with the highest score was 11.9 gr/dl and the lowest score was 10.7 gr/dl. While the average haemoglobin level of post test (after they were given red dragon fruit juice) was 12.36 gr/dl with the highest level 14.9 gr/dl and the lowest level 10.2 gr/dl. This proved that the average haemoglobin level before they were given red dragon fruit juice was actually lower than that of after they were given red dragon fruit juice (11.63 gr/dl < 12.36 gr/dl).

Based on the research results, it was known that the average haemoglobin level of the pre test and post test was different. The first group of young females given beet juice showed the *paired simple t-test* result was with *p-value*  $0.26 > 0.005$ . We can draw a conclusion that there is a difference between the average haemoglobin level of pre test and post test in the first young females group (after they were given beet juice). From the average haemoglobin level of pre test and post test in the second group (after they were given red dragon fruit juice), we could get *p-value*  $0.033 > 0.05$ . This showed that there was a difference of the average haemoglobin level of pre test and post test in the second group (after they were given red dragon fruit juice).

The independent analysis result was used to know whether there was a difference of the average haemoglobin level of pre test and post test between the the first and second group (after they were given beet juice and red dragon fruit juice). We obtained *p-value*  $0.515 > 0.05$  for the beet group and  $0.516 > 0.05$  for the dragon fruit group. This means that there is no difference increase in haemoglobin level of post test in both groups given beet juice and dragon fruit juice. We did not observe this so that the final result of the research biases because of several factors such as having breakfast habit, social economic status, and gender.

The hypothesis test with independent sample test was intended to know whether there was a difference of the average haemoglobin level of post test in both groups (after they were given beet juice and red dragon fruit juice). We obtained p-value  $0.515 > 0.05$  for the beet group and  $0.516 > 0.05$  for dragon fruit juice. This means that there was no increase in the level haemoglobin of post test in both groups after they were given beet juice and red dragon fruit juice at Muhammadiyah Al Kaustar Junior High School. Young females need energy, protein, and more other nutrients to grow. They need more iron than young males because it is needed to replace iron missing during menstrual period. Anaemia is a condition which there is deficiency of the capacity of oxygen carrier in blood. Fruits and vegetables containing vitamins and minerals function as antioxidants, phytoestrogen, and inflammation agent [12]. The deficiency of a number of vitamins and minerals can cause anaemia [13]. Girls who lack nutrients suffer from anaemia. Consuming tannin in the long term can affect iron status [14]. Most of young females have normal menstrual cycle [15]. [16] Consuming fruits containing vitamin C such as beet can increase the haemoglobin level and limits food intake which can be an inhibitor. Fe affects the optimization of Fe absorption in body. Consuming iron in 7 days will increase the haemoglobin level. Haemoglobin is a protein rich in iron. Protein plays an important part in transporting iron in body which leads to a problem if there is a deficiency of iron. The lower protein and iron intakes are, the lower the level of haemoglobin is [17]. The dragon fruit is a cactus of *Hyllocereus* and *Selenicereus*. Although it is a new kind of fruit in Indonesia, many people consume this fruit as a medicine for increasing stamina and blood cells.

#### 4. Conclusion

The consumption of beet juice and dragon fruit juice did not affect the haemoglobin level in young females because of several factors such as having breakfast habit good, social economic status, and gender. Research juice is an effective method to promote the consumption of fruits and vegetables as well as a balanced diet which offer to reduce high risks of various diseases.

#### Acknowledgment

Thank you to the Chairperson of College of Health Sciences Muhammadiyah Klaten for providing the opportunity to carry out this research.

#### Author Contributions

Presentation (E.W); Methodology (E.S); Data analysis (C.H.M)

#### References

- [1] Y. Wahyuni and R. Dewi, "Gangguan siklus menstruasi kaitannya dengan asupan zat gizi pada remaja vegetarian," *Indones. J. Nutr.*, vol. 6, no. 2, pp. 76–81, 2018.
- [2] F. Vitiasaridessy, "Kadar hemoglobin dengan kejadian dismenorea pada remaja putri," *J. Edu Heal.*, vol. 4, no. 2, 2014.
- [3] B. N. Nurul Qomariyah Rista Andaruni, "Efektifitas pemberian tablet zat besi (Fe), Vitamin C dan jus buah jambu biji terhadap peningkatan kadar hemoglobin (Hb) remaja putri di Universitas Muhammadiyah Mataram," *Midwifery J.*, vol. 3, no. 2, pp. 104–107, 2018.
- [4] E. A. M. Citta Cendani, "Asupan Mikronutrien, Kadar Hemoglobin dan Kesegaran Jasmani Remaja Putri," *Med. Med. Indones.*, vol. 45, no. 1, pp. 26–33, 2011.
- [5] D. Siallagan, P. D. Swamilaksita, and D. Angkasa, "Pengaruh asupan Fe, vitamin A, vitamin B12, dan vitamin C terhadap kadar hemoglobin pada remaja vegan," *J. Gizi Klin. Indones.*, vol. 13, no. 2, pp. 67–74, 2016.
- [6] F. A. Nugroho, D. Handayani, and Y. Apriani, "Asupan Protein Nabati Dan Kejadian Anemia," *J. Gizi Pangan*, vol. 10, no. November, pp. 165–170, 2015.

- [7] M. C. Jacobus and A. umboh Umboh, "Perbedaan kadar hemoglobin pada remaja gizi baik yang tinggal di pegunungan dengan yang tinggal di tepi pantai," *Jurna e-Clinic (eCI)*, vol. 4, no. 1, 2016.
- [8] H. Ashar, D. K. Mulyantoro, Y. D. Nurcahyani, and M. Khaerunnisa, "Anemia Pada Anak Sekolah Dasar Di Daerah Endemik Gaki," *MGMI*, vol. 7, no. 2, pp. 91–98, 2016.
- [9] D. Handayani, E. P. Pamungkasari, and E. S. Sulaeman, "Application of Path Analysis on Incidence of Anemia in Female Adolescents," *J. Gizi Pangan*, vol. 14, no. 1, pp. 37–44, 2019.
- [10] O. Article, B. H. Rashidi, H. M. Afzali, F. Haghollahi, G. Ardalan, and H. K. Jaliseh, "Evaluation of some diet habit and physical activity in adolescents in Tehran (2011)," *JQUMS*, vol. 22, no. 1, pp. 13–23, 2018.
- [11] A. Abay, H. W. Yalew, and A. Tariku, "Determinants of prenatal anemia in Ethiopia," pp. 1–10, 2017.
- [12] J. L. Slavin and B. Lloyd, "Health Bene fi ts of Fruits and Vegetables 1," pp. 506–516, 2012.
- [13] P. K. Goswami, S. Ghodgaonkar, and G. V. anagha raut Vishwakarma, "Contribution of medicinal plants in treatment of anaemia and as a haematinic," *J. Pharm. Pharm. Sci.*, vol. 4, no. 11, pp. 687–702, 2015.
- [14] N. M. Delimont, M. D. Haub, and B. L. Lindshield, "The Impact of Tannin Consumption on Iron Bioavailability and Status : A Narrative Review 1 , 2," *Curr. Dev. Nutr.*, pp. 1–12, 2012.
- [15] Z. Y. Saira Dars, Khashia Sayed, "Relationship of menstrual irregularities to BMI and nutritional status in adolescent girls," *J. Med. Sci.*, vol. 30, no. 1, 2014.
- [16] S. A. N. Eti Rimawati, Erna Kusumawati, Elviera Gamelia, Sumarah, "Food Supplement Intervention For Increasing Hemoglobin Level On Pregnant Women," *J. Ilmu Kesehat. Masy.*, vol. 9, no. November, pp. 161–170, 2018.
- [17] A. H. Al Rahmad, "Pengaruh Asupan Protein dan Zat Besi (Fe) terhadap Kadar Hemoglobin pada Wanita Bekerja," *J. Kesehat.*, vol. VIII, no. 3, pp. 321–325, 2017.
- [18] H. Arifin, Welli Nofiza, and Elisma, "Pengaruh Pemberian Jus Buah Naga *Hylocereus Undatus* (Haw.) Britt&Rose Terhadap Jumlah Hemoglobin, Eritrosit dan Hematokrit pada Mencit Putih Betina," *J. Sains dan Teknol. Farm.*, vol. 17, no. 2, pp. 118–125, 2012.