

## OUTHOR KORESPONDENSI

**Judul Artikel :** Study On Physical Distancing Compliance Towards Corona Virus Risk (COVID-19).

### Identitas Jurnal Ilmiah :

- a. Nama Jurnal: European Journal of Molecular & Clinical Medicine
- b. Nomor ISSN : ISSN – 2515-8260
- c. Volume, Nomor, Bulan Tahun : Volume 7, Issue 3, 2020
- d. Penerbit : EJCMC,
- e. DOI artikel (jika ada) : -
- f. Alamat Web Jurnal :  
<https://ejmcm.com/uploads/paper/418914357583e80f29698f4995b3b6c3.pdf>
- g. Terindeks di : Publon, Crossref, Embase DOAJ, Google scholar

### Lampiran Pendukung :

The screenshot displays the website for the European Journal of Molecular & Clinical Medicine (EJCMC). The page features a search bar at the top right with the text "Search for title, author, doi" and a "Search" button. Below the search bar, the journal's name "European Journal of Molecular & Clinical Medicine" is prominently displayed, along with the ISSN number "Online ISSN: 2515-8260". A navigation menu includes links for "About Journal", "Aims and Scope", "Editorial Board", "Archive", "Publication Ethics", "Peer Review Process", and "Contact Us". The main content area highlights the article "Study On Physical Distancing Compliance Towards Corona Virus Risk (COVID-19)" by Heni Sibawis Kejono<sup>1</sup>, Rizki Analia<sup>1</sup>, Nur Mustika Aji Nugroho<sup>2</sup>, and Setyono<sup>3</sup>. A "PDF" button is available for the article. The abstract text reads: "Abstract: The coronavirus (COVID-19) pandemic is a great concern in the global health sector. This led to the large number of people infected within a wide area. The possibility of transmission from one person to another increases the need for isolation or restrictions in terms of physical and social interactions within the community. This study aims to determine the extent of community compliance in conducting Physical Distancing. The research design was observational analytic, using cross-sectional method and was performed with 753 respondents. In addition, data was collected using internet-based questionnaire in the first Government Emergency Response period from March 23 to April 11, 2020. The information obtained includes demographics." A "Submit Your Article Here" button is also visible on the right side of the page. The bottom of the screenshot shows a Windows taskbar with the date and time "15:37 27/10/2023".

# Study On Physical Distancing Compliance Towards Corona Virus Risk (COVID-19).

Heru Subaris Kasjono<sup>1</sup>, Rizki Amalia<sup>1</sup>, Nur Mustika Aji Nugroho<sup>2</sup>, Sutaryono<sup>3</sup>

<sup>1</sup>Health Polytechnic, the Indonesian Health Ministry of Yogyakarta, Indonesia

<sup>2</sup>Health Literacy Foundation, Yogyakarta, Indonesia

<sup>3</sup>Muhammadiyah Health Sciences College of Klaten, Indonesia

e-mail : [heru.subarisk@poltekkesjogya.ac.id](mailto:heru.subarisk@poltekkesjogya.ac.id)

## Abstract

*The coronavirus (COVID-19) pandemic is a great concern in the global health sector. This is due to the large number of people infected within a wide area. The possibility of transmission from one person to another raises the need for isolation or restrictions in terms of physical and social interactions within the community. This study aims to determine the extent of community compliance in conducting Physical Distancing.*

*The research design was observational analytic, using cross sectional method, and was performed with 753 respondents. In addition, data was collected using internet-based questionnaire in the first Government Emergency Response period from March 23 to April 11, 2020. The information obtained includes demographics, physical social compliance variables and risk level. Therefore, qualitative analysis was performed to evaluate the frequency distribution, while the relationship between risk categories with Covid 19 cases was determined using chi square.*

*The results showed the majority of respondents performed outdoor activities (60.4%), although with good health protocol awareness. Also, most participants wear masks (75.6%), conduct good hand washing habit (86.6%), always use hand sanitizers (76.5) and maintain a distance from others (87%). Furthermore, the risk categories of respondents were low (95.2%), hence a relationship is established between Covid 19 cases and risk categories. This is reflected in physical and social distancing compliance ( $p < 0.00001$ ), classified in the moderate category, with an odd ratio value of 9.42 (CI: 4.45-19.97).*

*In conclusion, moderate physical and social distancing compliance poses a greater risk for covid19 cases, thus the need to increase discipline with Large Scale Social Restrictions (PSBB) implementation.*

**Keyword :** Covid-19, compliance , physical, social distancing, risk

## 1. INTRODUCTION

On December 31, 2019, Wuhan City Health Commission, Hubei province, China reported a pneumonia case with no known etiology, at Wuhan's Huanan Seafood Wholesale or fish and live animal markets. (1) On January 9, 2020, the Chinese CDC reported a novel coronavirus (2019-nCoV) as the cause, and the World Health Organization (WHO) announced the new name as Corona Virus Pandemic Disease (COVID-19) on February 11, 2020. To date, the statistics have elevated by 2,995,758 positive cases and 204,987 people deaths, in more than

213 countries, hence the depiction as a global threat. (2) Based on a report in April 7, 2020, the pandemic prevalence increased in Indonesia, with 2,738 positive, 204 recovered and 221 dead cases. (3) Furthermore, a report in April 29, 2020, indicated a further increase, to 9,771, 1,391 and 784, respectively. (3)

Coronavirus is a single positive, encapsulated, non-segmented RNA virus, and there are a total of four genera, including alpha, beta, delta and gamma forms (Huang *et al.* 2020; Fehr & Perlman 2015; Dawei Wang, *et al.* 2020). This pleomorphic microorganism possesses capsules, and is characterized by round or elliptical particles, with a diameter of about 50-200 nm. In addition, the structure is cube-like with protein S located on the surface. (Huang *et al.* 2020; Fehr & Perlman 2015; Dawei Wang, *et al.* 2020). The virus is sensitive to heat and is effectively deactivated by disinfectants containing chlorine, lipid solvents at a temperature of 56 °C for 30 minutes, ether, alcohol, peroxylacetic acid, non-ionic detergents, formalin, oxidizing agents and chloroform, while chlorhexidine is not effective. This infection dominantly occurs in winter, and under environmental conditions with relatively high humidity. (5,6)

This new Coronavirus pneumonia befalls mainly immunocompromised individuals. Despite the normal functioning of the body's immune system, high exposure to the virus at a time causes diseases. These manifestations progress faster and more severely in people with weak immune systems, including the elderly, pregnant women, and other conditions. In addition, the infection weakens the immune system against this virus, thus there is a tendency for re-infection. (5,6)

The main clinical symptoms of COVID-19 infection include fever (temperature > 38°C), cough and difficulty in breathing. This is possibly accompanied by heavy tightness, fatigue, myalgia, gastrointestinal symptoms, e.g., diarrhea and other respiratory symptoms. Furthermore, some patients tend to experience mild symptoms unaccompanied by fever. (2,5,6)

The spread of coronavirus is realized through close contact, the environment or contaminated objects, airway droplets, and airborne particles. Specifically, a droplet is a water-filled particle with a diameter of > 5 µm, and is capable of passing certain distance (usually 1 meter) to a vulnerable mucosal surface. The units are large enough, leading to poor ability to last or settle in air over a long period. These droplets originate from the airways, and are propagated by coughing, sneezing or talking, and also through invasive acts of respiratory procedures, including sputum aspiration, bronchoscopy, and tracheal tube insertion. In addition, airborne particles with diameter less than 5 µm have a potential to spread over long distances and remain infectious. Also, pathogens of this size spread by direct contact, or via blood and fluid into the body through mucous membranes or damaged skin. (6)

There are currently no vaccines available to ensure prevention, thus the best way to stop infection is by avoiding exposure to the virus. This is attainable through personal hygiene efforts on an individual level, by means of regular hand washing with soap, closing the mouth while coughing and ensuring the use of face masks, as well as increasing self-immunity and controlling comorbidities. Meanwhile, prevention at the community level involves the implementation of Large Scale Social Restrictions (PSBB), through physical (Physical Distancing) and social (Social Distancing) restrictions. (8)

The spread of COVID-19 is expected to be more extensive and faster on instances where there is limited participation and commitment from all parties, including the government, community and other sectors. Therefore, this study aims to assess compliance with Large-Scale Social Restrictions (PSBB) implementation created by the government, comprising the act of limiting physical (Social Distancing) and social (Social Distancing) interactions in the community. This research results are projected to be applied in evaluating and determining countermeasures for the next COVID-19 outbreak in Indonesia.

## 2. METHOD

The study design was observational analytic, using the cross sectional method, with 753 respondents. In addition, data was collected using an internet-based questionnaire, through the google doc application distributed by telephone and chat app. The responses were obtained according to the first Government Emergency Response Period, March 23 to April 11, 2020, including demographics, physical and social distancing compliance variables, as well as the risk level. Therefore, data analysis was performed qualitatively to evaluate the frequency distribution, while the relationship between risk categories and Covid 19 case was determined using chi square, with a significance level of 95%.

## 3. RESULTS

Research conducted using online media involved 753 respondents, with characteristic average age of 32 years, in a range between 15-75 years.

Based on the questionnaire, physical and social distancing compliance is shown in the following table:

Table. 1.1

*Descriptive analysis of physical and social distancing compliance*

| Activity  | Compliance |      |        |      |
|---|------------|------|--------|------|
|   | Yes (n)    | %    | No (n) | %    |
| Out of the house  | 455        | 60.4 | 298    | 39.6 |
| Online motorcycle ( <i>ojek</i> )/ taxi users                                       | 83         | 11.0 | 670    | 89.0 |
| City Transport Users  | 4          | 0.5  | 749    | 99.5 |
| Bus Users   | 10         | 1.3  | 743    | 98.7 |
| Bajaj User  | 0          | 0.0  | 753    | 100  |
| Train Users   | 7          | 0.9  | 746    | 99.1 |
| Using a mask to leave the house   | 569        | 75.6 | 184    | 24.4 |
| Doing a handshake   | 85         | 11.3 | 668    | 88.7 |
| Handwashing habits  | 652        | 86.6 | 101    | 13.4 |
| The habit of using a hand sanitizer   | 576        | 76.5 | 177    | 23.5 |
| Keep a distance from others when outside activities, study, work, worship, shopping | 655        | 87.0 | 98     | 13.0 |
| The washing hands habit after leaving the house                                     | 724        | 96.1 | 29     | 3.9  |
| Located in a contagious region / province   | 470        | 62.4 | 283    | 37.6 |
| Age 60 years and above  | 14         | 1.9  | 739    | 98.1 |

|  |     |      |     |      |
|--|-----|------|-----|------|
| History of heart disease, diabetes, chronic respiratory disorders  | 36  | 4.8  | 717 | 95.2 |
| There are ODP (People Under Monitoring) and PDP (Patients Under Surveillance) of Covid 19 in the respondent area | 207 | 27.5 | 546 | 72.5 |
| When working or studying at home, whether have a fever, runny nose, cough, or shortness of breath                | 95  | 12.6 | 658 | 87.4 |
| Travel history abroad or out of town (last 14-20 days)   | 103 | 13.7 | 650 | 86.3 |

Source: primary data

The results showed the participation of most respondents in outdoor activities (60.4%), withadequate awareness of health protocol. This is indicated by the majority wearing masks (75.6%), conductingregular hand washing habit (86.6%), always using hand sanitizers (76.5) and maintaining a distance from others (87%).

Based on the questionnaire,a high proportion of respondentswere in the low risk category (95.2%) for Covid 19 pandemic case (Table 1.2). Also, numerousparticipants live in the red zone area (62.4%), and are under 60 years old (98%). The presence of ODP (People Under Monitoring) and PDP (Patients Under Surveillance) was small (27%),withouta history of co-morbidity or concomitant diseases (95%). Therefore,a high proportion of respondents were indicated in the low risk category.

Table. 1.2

Descriptive analysis of Covid 19 transmission risk categories

| Risk     | n   | %     |
|----------|-----|-------|
| Moderate | 36  | 4.8   |
| Low      | 717 | 95.2  |
| Total    | 753 | 100.0 |

Source : primary data

The relationship between risk categories and Covid 19 cases isdepicted in the following table:

Table. 1.3

The relationship between Covid 19 cases with physical and social distancing compliance risk categories

| Case                              | Risk         |     |         |      |
|-----------------------------------|--------------|-----|---------|------|
|                                   | Moderate (n) | %   | Low (n) | %    |
| Covid_19                          | 4            | 8,5 | 43      | 91,5 |
| Suspect Covid_19                  | 0            | 0   | 7       | 100  |
| Patients Under Surveillance (PDP) | 0            | 0   | 9       | 100  |
| People Under Monitoring           | 12           | 25  | 36      | 75   |

| (ODP)       |    |      |     |      |
|-------------|----|------|-----|------|
| Do not know | 10 | 14,3 | 60  | 85,7 |
| Healthy     | 10 | 1,7  | 562 | 98,3 |

Source: Data processed

The analysis results showed p value  $<0.001$ , thus indicating the existence of a relationship between Covid 19 cases and risk categories, as a reflection towards physical and social distancing compliance.

The relationship between risk categories and Covid 19 case risk can be seen in the following table:

Table. 1.4

Relationship between risk categories and Covid risk 19

| Case          | Risk         |      |         |      | P value | OR   |
|---------------|--------------|------|---------|------|---------|------|
|               | Moderate (n) | %    | Low (n) | %    |         |      |
| Covid_19 risk | 26           | 18.4 | 155     | 85.6 | 0.00001 | 9.42 |
| Healthy       | 10           | 1.7  | 562     | 98.3 |         |      |

Source: Data processed

Table 1.4 is a cross table used to calculate the Covid 19 case risk, comparing positive respondents and those without diagnosis, to healthy individuals. The analysis results generated a p value  $<0.00001$ , thus indicating the presence of a relationship between Covid 19 and the risk categories. The odd ratio value determined was 9.42 (CI: 4.45-19.97), implying a risk of 9.42 times in the moderate category respondents, compared to those in the low risk category.

#### 4. DISCUSSION

This study showed a relationship between Covid 19 case and the risk category of being exposed or the physical and social distancing implementation ( $p = 0.00001$ ). In line with Mona's research, numerous opportunities for the spread of viruses are created in social networks without social and physical distancing practices, as well as vice versa (9). In addition, person-to-person transmission is reducible through public health efforts. (10) The WHO recommends distance maintenance, in order to avoid close contact, especially with health care workers, and also the avoidance of red zoned countries or regions. Furthermore, individuals with respiratory infections symptoms are expected to perform cough etiquette, by maintaining a distance, covering the cough and sneeze with tissue or disposable clothing, and practicing regular handwashing. Moreover, health care facilities are required to improve infection prevention standards and control practices, especially in emergency departments. (2) The Indonesian Health Ministry issued Regulation No. 9 of 2020 is based on Large-Scale Social Restriction (PSBB) Guidelines. Some limited activities include schooling, office work, religious activities, public facilities, social culture, public transportation as well as defense and security. This policy is, however, incomplete, as evidenced by the increasing case of Covid 19 in Indonesia based on the record of April 29, 2020, comprising 9,771 positive and 784 death incidents. (3) The upsurge was attributed to several factors, especially the level of

community compliance with the physical and social distancing implementation. The results showed the participation of most respondents in outdoor activities (60.4%), because of certain reasons, including the performance of designated tasks, working and others. The PSBB policy has not implemented a total lockdown, therefore allowing public freedom outdoors.

However, perpetrators have good awareness of health protocol, evidenced by wearing of face masks (75.6%), having the habit of handwashing (86.6%), regular use of hand sanitizers (76.5) and maintaining a distance from others (87 %). This results indicate the respondents to be in low risk category (95.2%), although most participants were below 60 years old (98%), and the presence of ODP (People Under Monitoring) as well as PDP (Patients Under Surveillance) was small (27%). Also, a majority presented with no history of comorbidity or concomitant diseases (95%), indicating a low risk of transmission.

Despite the efforts towards controlling Covid-19 worldwide, numerous unclear problems including the possibility of vaccines, transmission from animals to humans and inanimate objects, as well as the survival duration. Therefore, strict supervision and monitoring is needed for carriers without Symptoms (OTG), being a potential source of very dangerous infection to unsuspecting people. In addition, it is important for everyone to strictly implement health protocols, especially regular handwashing with disinfectants, maintaining a safe distance, and evading irrelevant outdoor activities. Furthermore, empowerment towards transmission prevention is achievable by involving all components of the community and exploring the local potential in each region.

The Odd ratio results of 9.42 showed the important role of physical and social distancing compliance in Covid 19 prevention. Therefore, respondents in the moderate category have a 9.42 times greater risk than the low group. Hence, greater observance of distancing rules reduces the tendency for infection. In Contagion theory, Corona virus spreads contagiously, and the term "contagion" refers to infections with the ability to spread rapidly in a network, including disasters and flu. This term was first used by Giralamo Fracastor in 1546. (9) Therefore, compliance to physical and social distancing are the most effective ways to control Covid 19 cases, although there are other protective efforts according to the WHO protocol. These include the implementation of five personal protective measures, which involve avoiding the touching of eyes, nose and mouth, alongside maintaining hand hygiene and keeping a distance. (11)

## 5. Conclusion

Communities with adequate awareness towards physical and social distancing compliance in the moderate category have a greater spread risk of Covid 19 cases. This prompts the need to increase discipline in implementing physical and social distancing compliance policies, by educating the people, especially in red zone areas.

## 6. REFERENCES

- [1] Lu, Hongzhou CWS, Tang and Y-W. Outbreak of Pneumonia of Unknown Etiology in Wuhan China: the Mystery and the Miracle. 2019;0-3.
- [2] WHO. (World Health Organization (WHO). Novel corona virus(2019-nCoV). Situation report. [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200428-sitrep-18-ncov.pdf?sfvrsn=fa644293\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200428-sitrep-18-ncov.pdf?sfvrsn=fa644293_2)).

- [3] Gugus Tugas Percepatan Penanganan COVID-19. Peta Sebaran Covid-19 [Internet]. 2020. p. diakses 29 april 2020. Available from: <https://covid19.go.id/peta-sebaran>
- [4] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Articles Clinical features of patients infected with 2019 novel coronavirus in Wuhan , China. 2020;6736(20):1–10.
- [5] Fehr AR, Perlman S. Chapter 1 Coronaviruses : An Overview of Their Replication and Pathogenesis. 2015;1282(1).
- [6] Dawei Wang, MD; Bo Hu, MD; Chang Hu, MD; Fangfang Zhu, MD; Xing Liu, MD; Jing Zhang, MD; Binbin Wang, MD; Hui Xiang M, Zhenshun Cheng, MD; Yong Xiong, MD; Yan Zhao, MD; Yirong Li, MD; Xinghuan Wang, MD; Zhiyong Peng M. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. 2020;1–9.
- [7] Erlina Burhan, Fathiyah Isbaniah, Agus Dwi Susanto , Tjandra Yoga Aditama, Soedarsono, Teguh Rahayu Sartono, Yani Jane Sugiri, Rezki Tantular, Bintang YM Sinaga , R.R Diah Handayani HA. Pneumonia Covid-19 Diagnosis & Penatalaksanaan Di Indonesia Tim. 1st ed. Jakarta: Perhimpunan Dokter Paru Indonesia; 2020.
- [8] Penanganan P, Virus C. Corona virus disease 2019. 2020;
- [9] No V, Mona N. Konsep Isolasi Dalam Jaringan Sosial Untuk Meminimalisasi Efek Contagious ( Kasus Penyebaran Virus Corona Di Indonesia ) Jurnal Sosial Humaniora Terapan. J Sos Hum Terap. 2020;2(2):117–25.
- [10] Song F, Shi N, Shan F, Zhang Z, Shen J, Lu H, et al. Emerging Coronavirus 2019-nCoV Pneumonia. Radiology. 2019;
- [11] Machida M, Nakamura I, Saito R, Nakaya T. International Journal of Infectious Diseases Adoption of personal protective measures by ordinary citizens during the COVID-19 outbreak in Japan. Int J Infect Dis [Internet]. 2020;94:139–44. Available from: <https://doi.org/10.1016/j.ijid.2020.04.014>